

REVIEW

Open Access



Regulation of mitochondrial functions by protein phosphorylation and dephosphorylation

Sangbin Lim¹, Kelly R. Smith¹, Ssang-Taek Steve Lim², Rong Tian³, Jianrong Lu⁴ and Ming Tan^{1,2*}

Abstract

The mitochondria are double membrane-bound organelles found in most eukaryotic cells. They generate most of the cell's energy supply of adenosine triphosphate (ATP). Protein phosphorylation and dephosphorylation are critical mechanisms in the regulation of cell signaling networks and are essential for almost all the cellular functions. For many decades, mitochondria were considered autonomous organelles merely functioning to generate energy for cells to survive and proliferate, and were thought to be independent of the cellular signaling networks. Consequently, phosphorylation and dephosphorylation processes of mitochondrial kinases and phosphatases were largely neglected. However, evidence accumulated in recent years on mitochondria-localized kinases/phosphatases has changed this longstanding view. Mitochondria are increasingly recognized as a hub for cell signaling, and many kinases and phosphatases have been reported to localize in mitochondria and play important functions. However, the strength of the evidence on mitochondrial localization and the activities of the reported kinases and phosphatases vary greatly, and the detailed mechanisms on how these kinases/phosphatases translocate to mitochondria, their subsequent function, and the physiological and pathological implications of their localization are still poorly understood. Here, we provide an updated perspective on the recent advancement in this area, with an emphasis on the implications of mitochondrial kinases/phosphatases in cancer and several other diseases.

Keywords: Mitochondria, Kinase, Phosphatase, Phosphorylation, Metabolism

Background

The mitochondria are double membrane-bound organelles found in most eukaryotic cells. They generate most of the cell's energy supply of adenosine triphosphate (ATP). Protein phosphorylation/dephosphorylation is a crucial regulatory system of signal transduction which controls many aspects of cellular functions. It can greatly impact the properties of a protein's enzymatic activity, structure, subcellular localization and stability. Phosphorylation, accomplished via a kinase action, and its reverse-de-phosphorylation via a phosphatase are essential switching-on and -off mechanisms in cell signaling. Many kinases and phosphatases have been

well-documented to play roles in the cell primarily outside mitochondria. Although the function and mechanisms of translocation to mitochondria are not well understood, the evolution of imaging and molecular techniques has allowed the discovery that an increasing number of kinases and phosphatases localize to mitochondria. The importance of phosphorylation/dephosphorylation in the governing of mitochondrial processes is supported by recent evidence that mitochondria contain many protein kinases, phosphatases, and associated proteins.

The aim of this review is to provide an updated perspective on the kinases and phosphatases localized to mitochondria. We describe here kinases and phosphatases that associate with mitochondria, their function within mitochondria, identified thus far, and the possible physiological or pathological implications of their mitochondrial localization.

*Correspondence: mtan@health.southalabama.edu

¹ Center for Cell Death and Metabolism, Mitchell Cancer Institute, University of South Alabama, Mobile, AL, USA

Full list of author information is available at the end of the article

Kinases localized in mitochondria

Tyrosine kinases

Abl

Abl is a non-receptor tyrosine kinase and is ubiquitously expressed in several cell types and tissues. *Abl* is localized at distinct sub-cellular sites including the nucleus, cytoplasm and endoplasmic reticulum (ER) and has various binding partners including signaling adaptors, proliferation regulators, transcription factors, kinases, phosphatases and cytoskeletal proteins [1]. While *Abl* is primarily localized to the nucleus, ER, and cytoplasm, it has been found that about 4 % of the total *Abl* resides in the mitochondria of murine embryo fibroblasts [2]. It has been revealed that H₂O₂ treatment of various cell types causes translocation of *Abl* to mitochondria and then induces cell death. Apart from H₂O₂, stimuli of ER stress, calcium ionophore A23187 and brefeldin A, also promote the translocation of *Abl* to mitochondria [3]. Although *Abl* does not contain a typical mitochondrial localization signal of its own, a study showed that protein kinase C δ binds to *Abl* in the ER. This PKC δ -*Abl* complex translocates from ER to mitochondria and then triggers apoptosis [4].

Abl contributes to human diseases such as Alzheimer's disease (AD) and chronic leukemia. Exposure of cultured neurons to various forms of multimeric amyloid- β peptide led to an increase in *Abl* tyrosine kinase activity and subsequent tyrosine phosphorylation of tau at Y394 [5–7]. Additionally, it was found that abnormal tau phosphorylation and subsequent cell death could be prevented by the *Abl* family kinase inhibitor [6, 7].

In chronic myeloid leukemia (CML) and acute promyelocytic leukemia (AML), *Abl* is activated upon translocation within breakpoint cluster region (Bcr) gene [8, 12]. Bcr/*Abl* fusion protein is a constitutively active tyrosine kinase which allows cells to proliferate without cytokine regulation, leading to a clonal myeloproliferative disorder. Its inhibitor, imatinib mesylate, can trigger apoptosis of CML cells [9]. It will be intriguing to explore if *Abl* or Bcr/*Abl*-associated apoptosis in AD and chronic leukemia is directed by mitochondrial localized *Abl* activity.

Src family kinases

There are multiple lines of evidence support the mitochondrial localization and function of *Src* family kinases. Although *Src* family kinases are permanent residents of cytoplasm, a previous study has suggested *Src* family including *Src*, *Fyn*, *Lyn*, *Fgr* and a negative regulator of *Src* family kinases, *CSK*, can be localized in mitochondria [10]. The phosphotyrosine signal in mitochondria was dramatically decreased by pretreatment with a *Src* inhibitor, PP2. This result suggests that *Src* family kinases may be associated with mitochondrial protein

phosphorylation. Other groups performed subfractionation of mitochondria using proteinase K and/or Triton X-100, further supported the localization of *Src* family kinases within mitochondria [11].

Since *Src* family kinases do not contain typical mitochondrial localization signals, they seem to be dependent on unidentified adaptor proteins for translocation into mitochondria. Several studies revealed that two anchoring proteins of protein kinase A (PKA), A Kinase Anchor Proteins 121 (AKAP121) and Dok-4, associate with *Src*. It is well known that AKAP121 acts as a multifunctional protein binding to protein tyrosine phosphatase D1 (PTPD1). PTPD1 anchors *Src* to the outer membrane of mitochondria upon activating the protein [12, 13]. Interestingly, a study showed that AKAP121 is found in the mitochondrial inner membrane with *Src* and *Lyn* [14]. Dok-4 is an adapter protein and appears to be responsible for the mitochondrial importation of *Src* in bovine endothelial cells. While deficiency of Dok-4 induces *Src* localization to extra-mitochondria, overexpression of Dok-4 causes *Src* localization to mitochondria [15].

Mitochondrial electron transport chain (ETC) is the final component of aerobic respiration, which contains a series of electron transporters embedded in the inner mitochondrial membrane. It includes complex I to IV and creates a chemical gradient that allows for the production of ATP. Several ETC complexes have been identified as substrates for *Src*. The first protein shown to be activated by mitochondrial *Src* was cytochrome c oxidase (complex IV in ETC) [11]. *Src* phosphorylates a subunit of cytochrome c oxidase, and leads to the enzymatic activation of complex IV in osteoclasts. Recently, it has been found that *Src* also affects other complexes within the ETC. One study showed that an increase in *Src* activity, as a response to changes in ATP levels in mitochondria of the rat brain, leads to an increase in the activities of complexes I, III, and IV, along with decreases in activity of complex V [16]. Two groups reported that inhibition of *Src* activity led to reduction in mitochondrial respiration mediated by a specific decrease in actions of complex I's NADH dehydrogenase-ubiquinone oxidoreductase system. NADH ubiquinone flavoprotein 2 at Tyr 193 of complex I and succinate dehydrogenase A at Tyr 215 of complex II, are the targets of *Src* phosphorylation [17, 18].

c-*Src*, normal proto-oncogene *Src*, possesses anti-apoptotic properties and shows increased protein levels and activity in a variety of human cancers [19]. Further, c-*Src* has been found to be a crucial player in multiple signaling pathways regulating proliferation, survival, metastasis, and angiogenesis [19]. One of these pathways is that of the mitochondrial, hypoxia-mediated ROS-induced activation of c-*Src*, HIF-1 α , and NF- κ B which

contributes to cell survival in various cancers including hepatomas, colon carcinomas, and neuroblastomas [20]. Additionally, c-Src has an important role in serum deprived pancreatic cancer cell survival [21]. Also, c-Src leads to metastasis of diverse cancers including breast cancer [22].

Mitochondrial Fgr kinase regulates complex II activity by tyrosine phosphorylation and plays an important role in regulating NADH/FADH₂ ratio. H₂O₂ promotes Fgr localization to mitochondria, but it does not seem that Fgr has a consensus signal sequence for mitochondrial localization [23].

Hibbs et al. found that mice homozygous for Lyn locus aberrations show abnormalities connected to B lymphocyte lineage and mast cell function [24]. Lyn^{-/-} mice did not mediate an allergic response to IgE cross-linking and exhibited severe glomerulonephritis caused by the kidney deposition of IgG immune complexes [24, 25], indicating that Lyn is associated with autoimmune disease. Lyn was also discovered to be critical for maintenance of the leukemic phenotypes of many different hematopoietic cancers including AML, CML and B cell lymphocytic leukemia [26–28]. Lyn was also expressed in some solid tumors. Therefore, it could serve as a potential therapeutic target for prostate cancer, glioblastomas, and aggressive subtypes of breast cancer [29–31]. Recent study revealed that Lyn-mediated mitochondrial tyrosine phosphorylation is required for hepatocyte survival under partial hepatectomy, pro-apoptotic conditions [32]. However, the role of Lyn in mitochondria and human diseases needs to be further elucidated.

Receptor tyrosine kinases

EGFR

A recent study demonstrated that epidermal growth factor receptor (EGFR) is localized to mitochondria where it interacts with cytochrome c oxidase subunit II (CoxII) in mouse fibroblasts over-expressing EGFR and Src [33]. EGF stimulation promotes translocation of EGFR into mitochondria. The translocation depends upon the phosphorylation of EGFR on Y845 by Src. The translocation is not observed upon overexpression of a catalytically inactive Src mutant or upon replacement of Y845 by phenylalanine (Y845-EGFR). A similar study showed that CoxII was phosphorylated by Src and EGFR, and the EGFR's mitochondrial localization was increased by EGF stimulation. Mitochondrial EGFR decreased the activity of Cox and reduced cellular ATP, indicating that mitochondrial EGFR regulates the function of mitochondria [34]. Moreover, new evidence showed that EGFR and EGFRVIII, a constitutively active variant of EGFR were localized in mitochondria [35, 36]. EGFR co-localized with FAK and Src in the mitochondria in glioma and xenografts. This

suggests a role of mitochondrial EGFR in cancer progression [37].

There is a debate as to whether the translocation of EGFR depends upon its endocytosis. One group concluded that the EGFR's localization to mitochondria depends on its endocytosis [38]. The report revealed that EGFR's mitochondrial translocation can be dramatically decreased in the presence of 3-methyladenine (3-MA), an autophagy inhibitor. Also, the knockdown of Beclin 1, an autophagy related gene, markedly decreased the mitochondrial translocation of EGFR. However, another group suggested that mitochondria-localized EGFR is independent of its internalization [39]. Therefore, the detailed mechanisms of EGFR mitochondrial localization still remain unclear.

ErbB2

ErbB2 is a human version of EGFR2. Recent study revealed that ErbB2 localizes to mitochondria of breast cancer cells and tumor samples of patients [40]. The study also found that localization of ErbB2 into mitochondria is mediated via association with mtHSP70 and reduces mitochondrial respiratory functions, including oxygen consumption, while it enhanced cellular glycolysis. It also showed that mitochondrial ErbB2 decreased membrane potential as well as the cellular ATP levels. However, detailed molecular mechanisms or direct substrates of mitochondrial EGFR and ErbB2 remain unclear. It is well known that EGFR and ErbB2 have been associated with the development of numerous human cancers. Tumors with changes in EGFRs tend to be more aggressive, and are considered as indicators of a poor clinical outcome, therefore they are intensely studied as therapeutic targets [41].

Serine/threonine kinases

Akt

It is well known that activated protein kinase B (Akt) is localized to diverse subcellular compartments: these include the Golgi, endoplasmic reticulum, and the nucleus. Also, it was shown that Akt could be localized in mitochondria when stimulated by 17 β -estradiol and insulin in endothelial cells [42]. In human neuroblastoma, it was reported that the activation of PI3 K/Akt signaling via insulin or insulin-like growth factor-1, is greatly amplified by mitochondrial Akt localization [43]. Recent studies also showed that treatment with leukemia inhibitory factor (LIF) increased the amount of total and phospho-Ser473-Akt in mitochondria [44], and insulin treatment could translocate Akt to mitochondria in cardiac muscle cells [45].

Several studies reported that mitochondrial Akt causes phospho-inactivation of the pro-apoptotic protein BAD,

and recruitment of Raf-1 to the mitochondria promoted cell survival [46, 47]. Recent studies also discovered new targets for mitochondrial Akt such as mitochondrial electron complex V and hexokinase-II [44, 45]. Akt isoform 1 modulates mitochondrial complex V activity, enhances the production of ATP, and increases phosphocreatine in cardiomyocytes, indicating that Akt is associated with ATP metabolism [45]. Akt also has a direct effect in mitochondria, which is mediated by phosphorylation of hexokinase-II, resulting in protection of the mitochondria from oxidant or Ca^{2+} -induced mitochondrial permeability transition pore (MPTP) opening [44]. Yang et al. found that translocation of phospho-Akt to mitochondria was enhanced in the streptozotocin-induced diabetic mice and insulin stimulates translocation of Akt to mitochondria [48]. Barksdale et al. revealed that HSP90 is responsible for Akt accumulation in mitochondria in unstimulated cells [49]. Taken together, mitochondrial Akt may play important roles in energy metabolism associated with diabetes.

JNK

It was well established that the c-Jun N-terminal kinase (JNK) localizes in cytoplasm and the nucleus. Many studies show that JNK may be activated in or translocated to the mitochondria. It was first revealed that JNK is localized in mitochondria in the phorbol ester response of myeloid leukemia cells [50]. Recent studies suggested an important role of JNK in mitochondria-related functions. These studies include murine heart mitochondria, hydrogen peroxide-treated rat brain or primary cortical cultures, acetaminophen-induced liver damage, multiple myeloma cells treated with anti-cancer drugs, and neonatal ischemia [51–57].

As mentioned before, JNK regulates apoptosis in general, indicating that targets of JNK can be apoptosis related proteins. JNK mediates phosphorylation and oligomerization of proapoptotic BAD, initiating apoptosis [58]. Conversely, there are studies reporting that JNK reduces apoptosis. For example, JNK is necessary for IL-3-mediated cell survival via phosphorylation and inactivation of BAD [59]. Also, activated JNK may colocalize with, and phosphorylate, Bcl-2 in mitochondrial membranes of hematopoietic cells [60]. This effect is present during interleukin-3-mediated stress, resulting in enhanced anti-apoptotic functions of Bcl-2. These contradictory results indicate that the localization and function of JNK in mitochondria are not well understood and warrant further investigation.

ERK1/2

Although extracellular signal-regulated kinase 1 and 2 (ERK1/2) normally translocates between cytosol and

the nucleus to influence trophic and pro-survival functions, recent studies found that ERK1/2 can localize to mitochondria of mouse heart [51], renal epithelial cells [61], mouse hippocampus [53] and human alveolar macrophages [62]. Mitochondrial ERK1/2 appears to play a crucial part in mitochondrial function [61, 62] including mitochondrial dysfunction, mitophagy, and apoptosis [63–65]. Recent investigation found that mitochondrial ERK1/2 phosphorylates steroidogenic acute regulatory protein (StAR), and this phosphorylation by ERK is required for the maintenance of this protein in mitochondria. Mitochondrial StAR together with mitochondrial active ERK and PKA are necessary for maximal steroid production [66].

p38 MAPK

Several studies revealed that the p38 mitogen-activated protein kinase (p38 MAPK) could reside in mitochondria. One group, using immunoblotting and immunofluorescence, showed that p38 MAPK is localized in cardiac mitochondria [51]. p38 MAPK was activated in the mitochondrial fraction by ischemia [67] and increased in mitochondria under H_2O_2 treatment [68]. Although it is well known that p38 MAPK performs a crucial function in mitochondria-mediated apoptosis, the precise function of mitochondrial p38 MAPK remains to be elucidated.

GSK3 β

It has been established that glycogen synthase kinase (GSK) 3 β can be localized in cytosol and nucleus. Some substrates of GSK3 β (e.g., tau), are cytosolic, whereas others (many transcription factors) are nuclear. However, in 1995, it was revealed that GSK3 β is present in mitochondria in rat cerebellum [69]. Since then several studies showed that GSK3 β regulates the function of mitochondrial proteins such as adenine nucleotide translocator (ANT) and cyclophilin D [70–72]. The mechanism of GSK3 β 's translocation to the mitochondria is unclear, but TOM20 may be integral, as in the example of connexin-43 [73]. In mitochondria, GSK3 β phosphorylates and suppresses a critical mitochondrial enzyme, pyruvate dehydrogenase activity [74]. Also, it has been demonstrated that GSK3 β has a role in determining the threshold for mitochondrial permeability transition pore (MPTP) opening [75], indicating that GSK3 β regulates mitochondria-mediated apoptosis.

Accumulated evidence shows that GSK3 β is highly associated with neurodegenerative diseases. GSK3 is a key kinase contributing to aberrant phosphorylation of the microtubule-binding protein tau in a process thought to cause neurofibrillary tangles in Alzheimer's disease [76]. In 1994, Mulot et al. found, via SDS-PAGE, that

paired helical filament-tau from the AD brain comprises four species which can be mimicked by GSK3 β mediated human brain tau phosphorylation in vitro [77]. Recent study also revealed that, in primary cortical neurons, GSK3 β is involved in Amyloid- β phosphorylation [78]. GSK3 β polymorphism was found to be associated with Parkinson's disease (PD) that alters GSK3 β transcription and splicing [79].

There is much interest in GSK3 β as a potential therapeutic target in many types of cancer. However, the application is complicated by findings that GSK3 β behaves as a tumor suppressor, but may promote cell proliferation in different types of cancer. There is evidence that GSK3 β functions as a tumor suppressor in skin and breast cancer [80, 81] and that repression of RNA polymerase 1 transcription by GSK3 β contributes to this tumor suppressor action [82]. However, there are contradictory findings that GSK3 β mediates tumor promotion and/or GSK3 β shows anti-proliferative effects in certain types of tumors including colon and pancreatic cancer [83, 84]. Whether these appeared contradictory functions of GSK3 β can be explained by the different localizations of this protein? It is still not clear if mitochondrial GSK3 β is associated with neurodegenerative diseases or cancer.

PKA

It has been acknowledged for many years that Protein kinase A (PKA) is crucial to mammalian mitochondrial physiology. Since the 1970s, it has been shown that PKA associates with, and in, mitochondria [85–88]. While catalytic subunits of PKA are predominantly found in the mitochondrial outer membrane [89], there is much evidence that PKA localizes to the mitochondrial inner membrane and matrix as well [87, 90–93]. In general, PKA is concentrated in cellular membranes and organelles through interactions with A kinase anchor proteins (AKAPs) [94, 95]. One group showed that de-localization of PKA from mitochondria was prompted by dominant negative AKAP121 mutation [96]. Indeed, expression of AKAP121 targets PKA to the cytoplasmic surface of mitochondria and then refines cAMP-PKA signaling to mitochondria [97]. Recent investigation revealed that AKAP1 recruits PKA and other signaling proteins to the outer mitochondrial membrane and thereby integrates several second messenger cascades by inhibitory phosphorylation of dynamin-related protein 1 (Drp1) and maintenance of mitochondrial integrity [98].

Drp1 is a mechanoenzyme using GTP hydrolysis to fuel the division of mitochondria. PKA mediated phosphorylation of Ser637 on Drp1 blocks Drp1's translocation to the mitochondrial surface and influences on fission [99]. Other findings indicate that outer mitochondrial PKA, and phosphatase PP2A, regulate neuronal development

by inhibiting and promoting mitochondrial division [100].

The cAMP response element-binding protein (CREB) is a transcription factor which regulates the transcription of cAMP response element-regulated genes and is found to bind to the mitochondrial DNA [101]. It has been shown that CREB is involved in regulation of mitochondrial gene expression and neuron longevity. Interestingly, it has been suggested that CREB may be phosphorylated by PKA within mitochondria [102].

Many heat shock proteins (Hsps) are known to play essential protective roles in the cardiovascular system. Hsp20 is expressed at high levels in cardiac, skeletal, and vascular smooth muscle. Hsp20 is regulated by the β -adrenergic/cAMP/PKA signaling pathway, which is known to be chronically activated in heart failure [103]. A recent study also revealed that PKA's phosphorylating Ser16 of Hsp20 is vital to the small Hsp's cardioprotective action [104]. For therapeutic development, it will be critical to distinguish if PKA mediated cardioprotective effects are due to mitochondrial PKA activity.

PKC

Protein kinase C (PKC) is a group of phospholipid-dependent serine/threonine kinases regulating numerous cellular functions through key signaling molecules. The activated PKCs translocate to multiple subcellular sites. Interestingly, analysis of the subcellular distribution of PKC ϵ in mouse heart with constitutively active PKC ϵ revealed that activated PKC ϵ is associated with a variety of mitochondrial proteins [51]. It has been suggested that mitochondrial translocation of PKC ϵ is associated with cardioprotection [105], and several studies showed that PKC ϵ substrates reside within cardiac mitochondria [106–108]. COX IV has been identified as a substrate of PKC ϵ in mitochondria [106]. Phosphorylation and activity of COX IV were increased by PKC activator, phorbol 12-myristate 13-acetate (PMA), but were blocked by the selective PKC ϵ inhibitor, ϵ V1-2. Another known mitochondrial target for PKC ϵ is the MPTP [107]. MPTP remains tightly closed under normal conditions, but it opens under cell death conditions. PKC ϵ interacts with presumed components of the MPTP in heart mitochondria, leading to reduced MPTP opening. Another target for PKC ϵ is a mitochondrial ATP-sensitive K⁺ channel (mitoK_{ATP}) [108]. It was recently found that PKC mediates interactions between connexin43 and mitoK_{ATP} subunit in mitochondria of cardiomyocytes [109]. These channels are normally closed, but are opened during periods of metabolic stress as ATP levels decline, leading to channel opening-mediated protective effects. This evidence reveals that mitochondrial PKC ϵ has a pro-survival role in a stress condition.

Hamasaki et al. found that expression of PKC ϵ decreases by 40 % in cardiac ventricles of hyperthyroid rats [110]. The expression of PKC ϵ was reduced in cytosol and membrane fractions but was not reduced in extracts of hypertrophied cardiac ventricles generated by aortocaval shunt or aortic banding. Other groups also revealed that hypertrophic stimuli activated PKC ϵ in rat-cultured cardiac myocytes, and in vivo, and that PKC ϵ overexpression in mice leads to cardiac hypertrophy related to concentric remodeling and the preservation of cardiac contractility [111, 112].

Recent study revealed that PKC ϵ increases the activity of endothelin converting enzyme, which degrades amyloid- β , and decreases amyloid plaque in transgenic mice [113], indicating that greater neuronal PKC ϵ activation may promote amyloid- β clearance, reducing Alzheimer's neuropathology. However, it has not been examined whether mitochondrial PKC ϵ is needed for AD progression.

PKC δ also translocates to mitochondria. Diverse studies revealed that treatment in a variety of neoplastic cells with hydrogen peroxide, phorbol esters, or anticancer agents such as cisplatin and etoposide causes PKC δ translocation into the mitochondria [114–116]. A study also found that mitochondrial PKC δ triggers the release of cytochrome c and resultant cell death. Thus, overexpressed PKC δ promotes apoptosis in neoplastic and normal keratinocytes by targeting mitochondria and disturbing their membrane potential [117]. However, direct targets for mitochondrial PKC δ still remain unclear.

PINK1

The PTEN-induced kinase 1 (PINK1) is a serine/threonine kinase. It is well known that PINK1 is a factor in autosomal recessive familial PD [118]. A large amount of evidence shows that PINK1 is localized in mitochondria and regulates mitochondrial function. A study found a strong mitochondrial targeting signal domain at its N terminus and showed that N-myc-tagged PINK1 expressed in mammalian cells accumulated in mitochondria. The protein could be found on the inner and outer mitochondrial membrane [119, 120]. PINK1 regulates Parkin (a protein related to PD), which acts as an E3 ubiquitin ligase [121]. PINK1 recruits Parkin to depolarized mitochondria. While PINK1 is imported and rapidly degraded by presenilin-associated rhomboid-like serine protease (PARL) in mitochondria with intact membrane potential, PINK1 breakdown is impaired in mitochondria with reduced membrane potential [120]. Apart from Parkin, PINK1 phosphorylates the mitochondrial fusion protein MFN2, which then acts as a mitochondrial receptor for Parkin [122] and Miro, an atypical Rho GTPase that tethers mitochondria to the tubulin network [123]. Several

studies showed that the PINK1/Parkin pathway is associated with fission and fusion events of mitochondria. Those studies revealed that the PINK1/Parkin in mitochondria regulates pro-fusion pathway given that the overexpression of PINK1 leads to interconnected and structurally elongated mitochondria, whereas knockdown of PINK1 leads to fragmentation of mitochondria [124, 125].

PINK1/Parkin pathway also regulates transport of mitochondria. Damaged mitochondria, prior to their clearance, are sequestered via the PINK1/Parkin pathway where mitochondrial movement is prevented through PINK1 mediated-Miro phosphorylation [123, 126]. PINK1 regulates mitochondrial content through mechanisms independent of mitophagy and PINK1-knockout mice showed a decreased Complex I activity [127, 128]. PINK1 mutant flies showed complex I deficiency and impaired ATP generation; however, supplementing these PINK1-deficient flies with complex 1 or ubiquinone corrects reductions in ATP generation allowing for recovery of flight muscle [129]. In both PINK1 and Parkin null flies, there lies a widespread defect with turnover of electron transport chain proteins [130]. Recent study demonstrated that PINK1 and Parkin mediate localized translation of respiratory chain component mRNAs along the mitochondrial outer membrane [131].

Dagda et al. studied the relationship between a PINK1 knock down and mitochondrial dysfunction in PD [132]. Stable knockdown of PINK1 produced mitochondrial fragmenting and autophagy in SH-SY5Y cells. Recent study found that loss of PINK1 accelerates neurodegenerative phenotypes induced by mitochondrial stress in transgenic animals where exists the conditional expression of mitochondrial unfolded ornithine transcarbamylase [133]. Mutations associated with PD were found throughout PINK1; however, the majority lay within the kinase domain, suggesting that loss of PINK1 kinase function is part of the pathogenesis [134]. Further, mutations within the carboxy terminus of PINK1 turn out to be critical to control an optimal kinase activity [135].

There is substantial evidence, based on epidemiological studies, for lower rates of cancer in patients with PD. The relationship between cancer incidence and PD may be related to the presence of genes common to both diseases [136]. Investigators found that PTEN knock-down suppresses PINK1 levels and shows cardioprotective effects, and concluded that PINK1 is a potential novel cardioprotective kinase [137]. However, the role of PINK1 in cancer development and progression is unclear. One study reported a link between Type II diabetes and PINK1 [138]. Investigators found that PINK1 was reduced in the skeletal muscle of Type II diabetic patients and in conditions of obesity or inactivity (or both) in patients and

controls. One study found glucose transporter inhibition under loss of PINK1 [139]. Recently, Deas et al. reported that the loss of PINK1 function seems to disturb glucose-sensing, resulting in enhanced insulin release, which is uncoupled from glucose uptake in beta cells [140], suggesting that a deficiency of the Parkinson's-associated PINK1 protein may directly alter insulin secretion at the pancreas.

Phosphatases translocate into mitochondria

Protein phosphatases

MKP1

MAP Kinase Phosphatases (MKPs) are dual specificity phosphatases which dephosphorylate both pSer/pThr and pTyr. MKP1 is localized in the nucleus through a LXXLL motif and appears to be confined to the dephosphorylation of nuclear MAPKs [141]. However, recent study found that MKP1 translocates to mitochondria after nerve growth factor treatment [142]. This result appears reasonable given that its substrates, and MAPKs, are also localized in mitochondria.

Wang et al. found that, in malignant samples, MKP1 is increased five fold over non-malignant samples [143]. This indicates that therapeutic suppression of MKP1 action may allow for expression of pro-apoptotic signaling of JNK in malignant cells. MKP1 is highly expressed in various human tumors, including non-small-cell lung cancer (NSCLC), bladder, ovarian, breast, osteosarcoma and prostate cancers [144–146]. They also found that MKP1 plays a critical part in the pathology of NSCLC, both in tumor growth and in response to treatment with cisplatin [146]. Recent study demonstrated that interference of MKP1 actions led to diminished invasion potential, tumor growth, and metastasis in mice [147].

A role for MKP1 in modulation of innate immune responses is strongly supported by several recent papers using models of endotoxic shock, anaphylaxis and arthritis. Mice deficient in MKP1 were hypersensitive to endotoxic shock; this was associated with prolonged activation of MAPKs, and enhanced production of TNF- α , IL-6, and IL-10 [148–150]. Moreover, it has been found that the absence of MKP1 markedly exacerbates disease development in mouse model of rheumatoid arthritis (RA). Overall, MKP1 controls dephosphorylation of MAPKs, but it is not clear whether this is the case for mitochondrial MAPKs and whether MKP1 is associated with human diseases due to mitochondrial MKP1 function.

Shp2

Src homology 2 domain-containing phosphatase 2 (Shp2) is ubiquitously expressed and contains two N-terminal SH2 domains along with a C-terminal protein tyrosine

phosphatase domain. Shp2 was primarily located inside the mitochondria related to cristae and the intercrystal space [16, 151]. Recent study also found that sepsis induced Shp2 mitochondrial localization and expression [152]. Mutations of Shp2 give rise to many distinct human diseases: Germ line Shp2 mutations result in Noonan Syndrome (NS), one of the most common autosomal dominant disorders, and LEOPARD Syndrome (LS), known, as its acronym reminds us, for its major manifestations: multiple Lentiginos, Electrocardiographic aberrations, Ocular hypertelorism, Pulmonary stenosis, Abnormalities of genitalia, Retardation of growth, and sensorineural Deafness [153, 154]. The distinct pathogenic routes of NS and LS are driven by different Shp2 mutations: NS-associated Shp2 mutations typically alter residues at the interface of N-SH2 and PTP domains [155], leading to increased enzymatic activity and RAS/ERK activation. LS mutations impact residues of the PTP domain, resulting in dramatic reduction of catalytic activity along with reduced activation of RAS/ERK [156].

Somatic Shp2 mutations are found in ~35 % of patients with sporadic juvenile myelomonocytic leukemia (JMML), a clonal pediatric myeloproliferative disorder (MPD) featuring the amplified expansion and tissue infiltration of myeloid cells, along with macrocytic anemia and persisting fetal hemoglobinemia [157, 158]. LS patients often undergo hypertrophic cardiomyopathy; in addition they may see an elevated potential for neuroblastoma and AML [159, 160]. Although shp2 expression and activity is closely associated with several diseases, connection between mitochondrial shp2 and disease progression has not been well established.

PTPs

Several protein tyrosine phosphatases (PTPs) such as PTPD1, PTP1B, and PTPMT1 has been reported as mitochondrial localized PTPs. PTPD1 is localized to outer membrane of mitochondria by binding to AKAP121 (or spliced isoform AKAP84) and increase a magnitude of EGF stimulated signaling [13]. In addition, PTPD1 binds to Src and activate Src, enhancing EGF-dependent mitogen signaling, enhancing oxidative phosphorylation and mitochondrial membrane potential [12]. PTP1B also localizes to mitochondria and enhances Src mediated mitochondrial oxidative phosphorylation [161]. Dual phosphatase PTPMT1 (PTP localized to the mitochondrion 1) has N-terminal mitochondrial localization signal sequence and is found in the matrix face. Knockdown of PTPMT1 alters mitochondrial phosphoprotein profile and markedly enhances ATP production [162]. This study clearly demonstrated the importance of protein phosphorylation/dephosphorylation switch in regulating mitochondrial function.

Lipid phosphatase


PTEN

Phosphatase and tensin homolog deleted on chromosome 10 (PTEN) is a well-known lipid phosphatase. Early studies proposed that PTEN localized exclusively to the cytoplasm and was able to transiently associate with the plasma membrane depending on the local PIP2 and PIP3 concentrations [163]. However, PTEN has recently been shown to localize to specialized subcellular compartments, such as the nucleus and the nucleolus, the mitochondria and the endoplasmic reticulum. Zhu et al. observed that a gradual buildup of PTEN in mitochondria occurred after induction of apoptosis, which was accompanied by the translocation of Bax to mitochondria [164]. Further, Zu et al. revealed that ischemia/reperfusion (I/R) induces mitochondrial localization of PTEN in the myocardium. Moreover, ischemic preconditioning attenuates mitochondrial localization of PTEN post-I/R, possibly blocking the translocation of Bax to the mitochondria, and leading to improved cell viability [165]. Liang et al. characterized an N-terminally extended form of PTEN (named PTEN α) that localizes to the cytoplasm and mitochondria, and induces cytochrome c oxidase activity and ATP generation in mitochondria [166]. This evidence showed that PTEN could exist in mitochondria, playing a crucial role in cellular functions such as apoptosis. However, whether this function of PTEN depends

on its lipid or protein phosphatase activities and what are the substrates in mitochondria are still not clear.

Molecular machinery for protein import into mitochondria

The majority of mitochondrial proteins are encoded in the nucleus, synthesized in the cytosol, and then delivered into their proper organelle. These mitochondrial proteins are imported into one of four mitochondrial compartments: outer membrane, intermembrane space, inner membrane, and matrix (Fig. 1). Each compartment contains translocases, which interact with precursor proteins to regulate their transport. The major molecular machinery which translocates proteins across the mitochondrial outer membrane is TOM (translocase of the outer membrane of mitochondria) complex. TOM comprises numerous integral membrane protein components: receptor subunits including Tom70 and Tom20, and core translocase subunits including Tom5, Tom6, Tom7, Tom22, and Tom40 [167, 168]. Upon transiting the channel of the TOM complex, substrate proteins may interact with one of three distinct machineries: translocase of inner mitochondrial membrane (TIM) complex 23, Tim9-Tim10 chaperone complex, and mitochondrial intermembrane space assembly machinery (MIA). The classic mitochondrial protein import pathway involves N-terminal presequences on the precursor proteins.



These proteins, following passage through the TOM complex, are directed to TIM23 complex and TIM23 bound presequence translocase-associated motor (PAM) which completes pre-protein translocation into the matrix. Following this, mitochondrial processing peptidases remove the presequences, and the proteins are folded to their native conformations. However, many mitochondrial precursor proteins are translated without cleavable presequences. These proteins also can be imported into mitochondria. The precursors of outer membrane beta-barrel proteins are transported, via the Tim9-Tim10 chaperone complex, to the sorting and assembly machinery (SAM) complex of the outer membrane [169]. Spanning proteins of the inner membrane also interact with the Tim9-Tim10 chaperone complex, after which they are inserted into the inner membrane by the TIM22 complex [170]. Several proteins of the intermembrane space contain cysteine motifs and are imported and oxidized by MIA [171].

As far, the detailed information of the mechanism of translocation of the kinases and phosphatases to mitochondria are elusive. The majority of the reported mitochondrial localized kinases/phosphatases lack typical mitochondrial localization signal. This raises three possibilities. First, it does not need a typical mitochondrial localization signal to be localized in mitochondria; second, there are hidden undiscovered mitochondrial localization signal; third, it is passively transported to

mitochondria by a cargo carrier. Since it has been shown that mitochondrial kinases/phosphatases play important roles in regulating various cellular functions, more investigations on the mechanism of the importation are definitely needed.


Concluding remarks and future directions

This review summarizes recent findings concerning mitochondrial localization of kinases and phosphatases, recaps our knowledge on their function to cellular domains beyond those commonly cited in textbooks (Table 1). Over the past many years, the role of mitochondria has been underestimated and understudied. Evidence revealed by recent advanced laboratory techniques such as tandem mass spectrometry, confocal imaging, and electron microscopy, reevaluate mitochondria as a crucial platform for cellular signaling beyond just energy production in the cell. Mitochondria have diverse reported phosphoproteins, kinases and phosphatases, as we have described here. Mounting evidence support that phosphorylation or dephosphorylation of mitochondrial proteins influences mitochondrial function, including metabolism of sugar, amino acids, and lipids; oxidative phosphorylation, antioxidant protein expression, mitochondrial fission and fusion, and decision for survival/death. Emerging evidence has shown that mitochondrial kinases/phosphatases regulate a variety of key regulatory processes in diverse mammalian

Table 1 Mitochondrial kinases/phosphatases

Proteins	Evidences of mitochondria localization	Functions in mitochondria	References
Abl	Co-IP, WB, IF	Apoptosis	[3, 4]
Src	WB, IP, Co-IP, Kinase array	Cytochrome C oxidase, ROS production and respiration	[10, 11, 15–18]
EGFR	WB, IP, IF	Apoptosis, respiration and cellular metabolism	[33–40]
Akt	WB, IP, Co-IP, IF	Cell survival and regulation of respiration	[43–45, 47–49]
JNK	WB, IF	Apoptosis, neuro inflammation and mitochondria biogenesis Possibly regulation of signal transduction	[50–57, 60]
ERK1/2	WB, IF	Respiration, ATP production, membrane potential, mitophagy, autophagic cell death, MPTP and steroidogenesis	[51, 53, 61–66]
P38 MAPK	WB, IF, GST	Apoptosis	[51, 67, 68]
GSK3 β	WB, Co-IP	MPTP and energy metabolism	[70–72, 74, 75]
PKA	WB, IF, DEAE, GST, in vitro kinase assay	Respiration, mitochondria division and steroidogenesis	[87, 89–93, 95, 98, 99, 102]
PKC	WB, IP, IF	Respiration, K ⁺ channel and apoptosis	[105–109]
PINK1	WB, IF	Mitochondria trafficking at outer membrane. Pathogenesis of PD. Localized translation of respiration chain complex	[121, 122, 125–128, 130, 131]
MKP1	WB, IF	Apoptosis	[142]
Shp2	WB, Co-IP	ROS production	[16, 151, 152]
PTPs	WB, Co-IP, IF	ATP production	[12, 13, 161, 162]
PTEN	WB, IF	Apoptosis and ATP production	[164, 165]

WB western blot, Co-IP Co-immunoprecipitation, IP Immunoprecipitation, IF Immunofluorescence, GST GST pull-down assay, DEAE diethylaminoethyl cellulose chromatography



tissue [172–175]. Intriguingly, a recent phosphoproteome analysis of functional mitochondria showed there are 155 phosphorylation sites in 77 mitochondrial proteins including inner membrane ETS and enzymes in resting human muscle [176]. Although the detailed functions of these kinases/phosphatases in mitochondria are still unclear, there is evidence that the translocated mitochondrial kinases/phosphatases may be seen in mitochondria in several mitochondrial compartments: Those in the matrix likely regulate enzymes of the TCA cycle, amino acid metabolism, free-radical balance, ETC activity, and ATP synthesis (Fig. 2). Those in the outer membrane regulate transport, cell death-related proteins, and mitochondrial fission/fusion. Those at the inner membrane regulate free-radical balance, nucleotide transport, ETC activity and complex assembly.

In summary, the research community has accumulated a large amount of evidence to support that phosphorylation and dephosphorylation of mitochondrial proteins influence mitochondrial function. However, the strength of the evidence on mitochondrial localization and their activities of the reported kinases and phosphatases vary greatly, and the detailed mechanisms on how these kinases/phosphatases translocate to mitochondria and the physiological and pathological roles related to their mitochondrial localization are still poorly understood. Due to increasing evidence supporting the important functions of mitochondrial kinases/phosphatases, mitochondrial biology is due for more intense exploration in this area.

Abbreviations

ER: endoplasmic reticulum; AD: Alzheimer's disease; CML: chronic myeloid leukemia; BCR: breakpoint cluster region; AML: acute promyelocytic leukemia; PKA: protein kinase A; AKAP121: a kinase anchor proteins 121; PTPD1: protein tyrosine phosphatase D1; ETC: mitochondrial electron transport chain; EGFRs: epidermal growth factor receptors; CoxII: cytochrome c oxidase subunit II; 3-MA: 3-methyladenine; SH: secondary hyper-parathyroidism; Akt: protein kinase B; LIF: leukemia inhibitory factor; MPTP: mitochondrial permeability transition pore; JNK: c-Jun N-terminal kinase; ERK1/2: extracellular signal-regulated kinase 1 and 2; p38 MAPKs: p38 mitogen-activated protein kinases; GSK: glycogen synthase kinase ANT: adenine nucleotide translocator; HD: Huntington's disease; Drp1: dynamin-related protein 1; CREB: cAMP response element-binding protein; DS: down syndrome; Hsps: heat shock proteins; PKC: protein kinase C; mitoK_{ATP}: mitochondrial ATP-sensitive K⁺ channel; PINK1: PTEN-induced kinase 1; PD: Parkinson's disease; PARL: presenilin-associated rhomboid-like serine protease; MKPs: MAP Kinase Phosphatases; NSCLC: non-small-cell lung cancer; RA: rheumatoid arthritis; Shp2: Src homology 2 domain-containing phosphatase 2; NS: Noonan syndrome; LS: Leopard syndrome; JMML: juvenile myelomonocytic leukemia; MPD: myeloproliferative disorder; PTP: protein tyrosine phosphatases; PTEN: phosphatase and tensin homolog deleted on chromosome 10; I/R: ischemia/reperfusion; TOM: translocase of the outer membrane of mitochondria; TIM: translocase of inner mitochondrial membrane; MIA: mitochondrial intermembrane space assembly machinery; PAM: presequence translocase-associated motor; SAM: sorting and assembly machinery.

Authors' contributions

SL and TM designed and wrote the manuscript, SL and KS collected and analyzed data; SL, STL, RT, JL and MT revised the manuscript. All authors read and approved the final manuscript.

Author details

¹ Center for Cell Death and Metabolism, Mitchell Cancer Institute, University of South Alabama, Mobile, AL, USA. ² Department of Biochemistry and Molecular Biology, College of Medicine, University of South Alabama, Mobile, AL, USA. ³ Mitochondria and Metabolism Center, Department of Anesthesiology and Pain Medicine, University of Washington, Seattle, WA, USA. ⁴ Department of Biochemistry and Molecular Biology, University of Florida College of Medicine, Gainesville, FL, USA.

Acknowledgements

We are Grateful to the support from The Vincent F. Kilborn, Jr. Cancer Research Foundation, National Institutes of Health Grants R01CA149646 (M. Tan), and American heart association scientist development Grant 12SDG10970000 (S.T. Lim).

Competing interests

The authors declare that they have no competing interests.

Received: 4 March 2016 Accepted: 1 April 2016

Published online: 14 April 2016

References

- Hantschel O, Superti-Furga G. Regulation of the c-Abl and Bcr-Abl tyrosine kinases. *Nat Rev Mol Cell Biol*. 2004;5(1):33–44.
- Kumar S, Bharti A, Mishra NC, Raina D, Kharbanda S, Saxena S, Kufe D. Targeting of the c-Abl tyrosine kinase to mitochondria in the necrotic cell death response to oxidative stress. *J Biochem*. 2001;276(20):17281–5.
- Ito Y, Pandey P, Mishra N, Kumar S, Narula N, Kharbanda S, Saxena S, Kufe D. Targeting of the c-Abl tyrosine kinase to mitochondria in endoplasmic reticulum stress-induced apoptosis. *Mol Cell Biol*. 2001;21(18):6233–42.
- Qi X, Mochly-Rosen D. The PKCdelta-Abl complex communicates ER stress to the mitochondria - an essential step in subsequent apoptosis. *J Cell Sci*. 2008;121(Pt 6):804–13.
- Derkinderen P, Scales TM, Hanger DP, Leung KY, Byers HL, Ward MA, Lenz C, Price C, Bird IN, Perera T, et al. Tyrosine 394 is phosphorylated in Alzheimer's paired helical filament tau and in fetal tau with c-Abl as the candidate tyrosine kinase. *J Neurosci*. 2005;25(28):6584–93.
- Williamson R, Scales T, Clark BR, Gibb G, Reynolds CH, Kellie S, Bird IN, Varndell IM, Sheppard PW, Everall I, et al. Rapid tyrosine phosphorylation of neuronal proteins including tau and focal adhesion kinase in response to amyloid-beta peptide exposure: involvement of Src family protein kinases. *J Neurosci*. 2002;22(1):10–20.
- Alvarez AR, Sandoval PC, Leal NR, Castro PU, Kosik KS. Activation of the neuronal c-Abl tyrosine kinase by amyloid-beta-peptide and reactive oxygen species. *Neurobiol Dis*. 2004;17(2):326–36.
- Ben-Neriah Y, Daley GQ, Mes-Masson AM, Witte ON, Baltimore D. The chronic myelogenous leukemia-specific P210 protein is the product of the bcr/abl hybrid gene. *Science*. 1986;233(4760):212–4.
- Holtz MS, Slovak ML, Zhang F, Sawyers CL, Forman SJ, Bhatia R. Imatinib mesylate (ST1571) inhibits growth of primitive malignant progenitors in chronic myelogenous leukemia through reversal of abnormally increased proliferation. *Blood*. 2002;99(10):3792–800.
- Salvi M, Brunati AM, Bordin L, La Rocca N, Clari G, Toninello A. Characterization and location of Src-dependent tyrosine phosphorylation in rat brain mitochondria. *Biochim Biophys Acta*. 2002;1589(2):181–95.
- Miyazaki T, Neff L, Tanaka S, Horne WC, Baron R. Regulation of cytochrome c oxidase activity by c-Src in osteoclasts. *J Cell Biol*. 2003;160(5):709–18.
- Livigni A, Scorziello A, Agnese S, Adornetto A, Carlucci A, Garbi C, Castaldo I, Annunziato L, Avvedimento EV, Feliciello A. Mitochondrial AKAP121 links cAMP and Src signaling to oxidative metabolism. *Mol Biol Cell*. 2006;17(1):263–71.
- Cardone L, Carlucci A, Affaitati A, Livigni A, DeCristofaro T, Garbi C, Varone S, Ullrich A, Gottesman ME, Avvedimento EV, et al. Mitochondrial AKAP121 binds and targets protein tyrosine phosphatase D1, a novel positive regulator of src signaling. *Mol Cell Biol*. 2004;24(11):4613–26.
- Sardanelli AM, Signorile A, Nuzzi R, Rasmussen DD, Technikova-Dobrova Z, Drahotova Z, Occhiello A, Pica A, Papa S. Occurrence of A-kinase anchor protein and associated cAMP-dependent protein kinase in the inner compartment of mammalian mitochondria. *FEBS Lett*. 2006;580(24):5690–6.
- Itoh S, Lemay S, Osawa M, Che W, Duan Y, Tompkins A, Brookes PS, Sheu SS, Abe J. Mitochondrial Dok-4 recruits Src kinase and regulates NF-kappaB activation in endothelial cells. *J Biochem*. 2005;280(28):26383–96.
- Arachiche A, Augereau O, Decossas M, Pertuiset C, Gontier E, Letellier T, Dachary-Prigent J. Localization of PTP-1B, SHP-2, and Src exclusively in rat brain mitochondria and functional consequences. *J Biochem*. 2008;283(36):24406–11.
- Hebert-Chatelain E, Jose C, Cortes NG, Dupuy JW, Rocher C, Dachary-Prigent J, Letellier T. Preservation of NADH ubiquinone-oxidoreductase activity by Src kinase-mediated phosphorylation of NDUFB10. *Biochim Biophys Acta*. 2012;1817(5):718–25.
- Ogura M, Yamaki J, Homma MK, Homma Y. Mitochondrial c-Src regulates cell survival through phosphorylation of respiratory chain components. *Biochem J*. 2012;447(2):281–9.
- Ishizawa R, Parsons SJ. c-Src and cooperating partners in human cancer. *Cancer Cell*. 2004;6(3):209–14.
- Lluis JM, Buricchi F, Chiarugi P, Morales A, Fernandez-Checa JC. Dual role of mitochondrial reactive oxygen species in hypoxia signaling: activation of nuclear factor-kappaB via c-SRC and oxidant-dependent cell death. *Cancer Res*. 2007;67(15):7368–77.
- Tejedo JR, Ramirez R, Cahuana GM, Rincon P, Sobrino F, Bedoya FJ. Evidence for involvement of c-Src in the anti-apoptotic action of nitric oxide in serum-deprived R1Nm5F cells. *Cell Signal*. 2001;13(11):809–17.
- Myoui A, Nishimura R, Williams PJ, Hiraga T, Tamura D, Michigami T, Mundy GR, Yoneda T. C-SRC tyrosine kinase activity is associated with tumor colonization in bone and lung in an animal model of human breast cancer metastasis. *Cancer Res*. 2003;63(16):5028–33.
- Acin-Perez R, Carrascoso I, Baixauli F, Roche-Molina M, Latorre-Pellicer A, Fernandez-Silva P, Mittelbrunn M, Sanchez-Madrid F, Perez-Martos A, Lowell CA, et al. ROS-triggered phosphorylation of complex II by Fgr kinase regulates cellular adaptation to fuel use. *Cell Metab*. 2014;19(6):1020–33.
- Hibbs ML, Tarlinton DM, Armes J, Grail D, Hodgson G, Maglito R, Stacker SA, Dunn AR. Multiple defects in the immune system of Lyn-deficient mice, culminating in autoimmune disease. *Cell*. 1995;83(2):301–11.
- Oracki SA, Tsantikos E, Quilici C, Light A, Schmidt T, Lew AM, Martin JE, Smith KG, Hibbs ML, Tarlinton DM. CTLA4lg alters the course of autoimmune disease development in Lyn-/- mice. *J Immunol*. 2010;184(2):757–63.
- Roginskaya V, Zuo S, Caudell E, Nambudiri G, Kraker AJ, Corey SJ. Therapeutic targeting of Src-kinase Lyn in myeloid leukemic cell growth. *Leukemia*. 1999;13(6):855–61.
- Donato NJ, Wu JY, Stapley J, Gallick G, Lin H, Arlinghaus R, Talpaz M. BCR-ABL independence and LYN kinase overexpression in chronic myelogenous leukemia cells selected for resistance to ST1571. *Blood*. 2003;101(2):690–8.
- Contri A, Brunati AM, Trentin L, Cabrelle A, Miorin M, Cesaro L, Pinna LA, Zambello R, Semenzato G, Donella-Deana A. Chronic lymphocytic leukemia B cells contain anomalous Lyn tyrosine kinase, a putative contribution to defective apoptosis. *J Clin Invest*. 2005;115(2):369–78.
- Goldenberg-Furmanov M, Stein I, Pikarsky E, Rubin H, Kasem S, Wygoda M, Weinstein I, Reuveni H, Ben-Sasson SA. Lyn is a target gene for prostate cancer: sequence-based inhibition induces regression of human tumor xenografts. *Cancer Res*. 2004;64(3):1058–66.
- Choi YL, Bocanegra M, Kwon MJ, Shin YK, Nam SJ, Yang JH, Kao J, Godwin AK, Pollack JR. LYN is a mediator of epithelial-mesenchymal transition and a target of dasatinib in breast cancer. *Cancer Res*. 2010;70(6):2296–306.
- Stettner MR, Wang W, Nabors LB, Bharasa S, Flynn DC, Grammer JR, Gillespie GY, Gladson CL. Lyn kinase activity is the predominant cellular SRC kinase activity in glioblastoma tumor cells. *Cancer Res*. 2005;65(13):5535–43.
- Gringeri E, Carraro A, Tibaldi E, D'Amico FE, Mancon M, Toninello A, Pagano MA, Vio C, Cillo U, Brunati AM. Lyn-mediated mitochondrial tyrosine phosphorylation is required to preserve mitochondrial integrity in early liver regeneration. *Biochem J*. 2010;425(2):401–12.
- Boerner JL, Demory ML, Silva C, Parsons SJ. Phosphorylation of Y845 on the epidermal growth factor receptor mediates binding to the mitochondrial protein cytochrome c oxidase subunit II. *Mol Cell Biol*. 2004;24(16):7059–71.
- Demory ML, Boerner JL, Davidson R, Faust W, Miyake T, Lee I, Huttemann M, Douglas R, Haddad G, Parsons SJ. Epidermal growth factor receptor translocation to the mitochondria: regulation and effect. *J Biochem*. 2009;284(52):36592–604.
- Cao X, Zhu H, Ali-Osman F, Lo HW. EGFR and EGFRvIII undergo stress- and EGFR kinase inhibitor-induced mitochondrial translocation: a

- potential mechanism of EGFR-driven antagonism of apoptosis. *Mol Cancer*. 2011;10:26.
36. Dreier A, Barth S, Goswami A, Weis J. Cetuximab induces mitochondrial translocation of EGFRvIII, but not EGFR: involvement of mitochondria in tumor drug resistance? *Tumour Biol*. 2012;33(1):85–94.
 37. Dasari VR, Velpula KK, Alapati K, Gujrati M, Tsung AJ. Cord blood stem cells inhibit epidermal growth factor receptor translocation to mitochondria in glioblastoma. *PLoS One*. 2012;7(2):e31884.
 38. Yue X, Song W, Zhang W, Chen L, Xi Z, Xin Z, Jiang X. Mitochondrially localized EGFR is subjected to autophagic regulation and implicated in cell survival. *Autophagy*. 2008;4(5):641–9.
 39. Yao Y, Wang G, Li Z, Yan B, Guo Y, Jiang X, Xi Z. Mitochondrially localized EGFR is independent of its endocytosis and associates with cell viability. *Acta Biochim Biophys Sin*. 2010;42(11):763–70.
 40. Ding Y, Liu Z, Desai S, Zhao Y, Liu H, Pannell LK, Yi H, Wright ER, Owen LB, Dean-Colomb W, et al. Receptor tyrosine kinase ErbB2 translocates into mitochondria and regulates cellular metabolism. *Nat Commun*. 2012;3:1271.
 41. Holbro T, Hynes NE. ErbB receptors: directing key signaling networks throughout life. *Ann Rev Pharmacol Toxicol*. 2004;44:195–217.
 42. Sasaki K, Sato M, Umezawa Y. Fluorescent indicators for Akt/protein kinase B and dynamics of Akt activity visualized in living cells. *J Biochem*. 2003;278(33):30945–51.
 43. Bijur GN, Jope RS. Rapid accumulation of Akt in mitochondria following phosphatidylinositol 3-kinase activation. *J Neurochem*. 2003;87(6):1427–35.
 44. Miyamoto S, Murphy AN, Brown JH. Akt mediates mitochondrial protection in cardiomyocytes through phosphorylation of mitochondrial hexokinase-II. *Cell Death Differ*. 2008;15(3):521–9.
 45. Yang JY, Deng W, Chen Y, Fan W, Baldwin KM, Jope RS, Wallace DC, Wang PH. Impaired translocation and activation of mitochondrial Akt1 mitigated mitochondrial oxidative phosphorylation complex V activity in diabetic myocardium. *J Mol Cell Cardiol*. 2013;59:167–75.
 46. Datta SR, Dudek H, Tao X, Masters S, Fu H, Gotoh Y, Greenberg ME. Akt phosphorylation of BAD couples survival signals to the cell-intrinsic death machinery. *Cell*. 1997;91(2):231–41.
 47. Majewski M, Nieborowska-Skorska M, Salomoni P, Slupianek A, Reiss K, Trotta R, Calabretta B, Skorski T. Activation of mitochondrial Raf-1 is involved in the antiapoptotic effects of Akt. *Cancer Res*. 1999;59(12):2815–9.
 48. Yang JY, Yeh HY, Lin K, Wang PH. Insulin stimulates Akt translocation to mitochondria: implications on dysregulation of mitochondrial oxidative phosphorylation in diabetic myocardium. *J Mol Cell Cardiol*. 2009;46(6):919–26.
 49. Barksdale KA, Bijur GN. The basal flux of Akt in the mitochondria is mediated by heat shock protein 90. *J Neurochem*. 2009;108(5):1289–99.
 50. Ito Y, Mishra NC, Yoshida K, Kharbanda S, Saxena S, Kufe D. Mitochondrial targeting of JNK/SAPK in the phorbol ester response of myeloid leukemia cells. *Cell Death Differ*. 2001;8(8):794–800.
 51. Baines CP, Zhang J, Wang GW, Zheng YT, Xiu JX, Cardwell EM, Bolli R, Ping P. Mitochondrial PKCepsilon and MAPK form signaling modules in the murine heart: enhanced mitochondrial PKCepsilon-MAPK interactions and differential MAPK activation in PKCepsilon-induced cardioprotection. *Circ Res*. 2002;90(4):390–7.
 52. Chauhan D, Li G, Hideshima T, Podar K, Mitsiades C, Mitsiades N, Munshi N, Kharbanda S, Anderson KC. JNK-dependent release of mitochondrial protein, Smac, during apoptosis in multiple myeloma (MM) cells. *J Biochem*. 2003;278(20):17593–6.
 53. Rumora L, Lovric J, Sairam MR, Maysinger D. Impairments of heat shock protein expression and MAPK translocation in the central nervous system of follitropin receptor knockout mice. *Exp Gerontol*. 2007;42(7):619–28.
 54. Hanawa N, Shinohara M, Saberi B, Gaarde WA, Han D, Kaplowitz N. Role of JNK translocation to mitochondria leading to inhibition of mitochondrial bioenergetics in acetaminophen-induced liver injury. *J Biochem*. 2008;283(20):13565–77.
 55. Zhou Q, Lam PY, Han D, Cadenas E. c-Jun N-terminal kinase regulates mitochondrial bioenergetics by modulating pyruvate dehydrogenase activity in primary cortical neurons. *J Neurochem*. 2008;104(2):325–35.
 56. Nijboer CH, Bonestroo HJ, Zijlstra J, Kavelaars A, Heijnen CJ. Mitochondrial JNK phosphorylation as a novel therapeutic target to inhibit neuroinflammation and apoptosis after neonatal ischemic brain damage. *Neurobiol Dis*. 2013;54:432–44.
 57. Chambers JW, Howard S, LoGrasso PV. Blocking c-Jun N-terminal kinase (JNK) translocation to the mitochondria prevents 6-hydroxydopamine-induced toxicity in vitro and in vivo. *J Biochem*. 2013;288(2):1079–87.
 58. Bhakar AL, Howell JL, Paul CE, Salehi AH, Becker EB, Said F, Bonni A, Barker PA. Apoptosis induced by p75NTR overexpression requires Jun kinase-dependent phosphorylation of Bad. *J Neurosci*. 2003;23(36):11373–81.
 59. Yu C, Minemoto Y, Zhang J, Liu J, Tang F, Bui TN, Xiang J, Lin A. JNK suppresses apoptosis via phosphorylation of the proapoptotic Bcl-2 family protein BAD. *Mol Cell*. 2004;13(3):329–40.
 60. Deng X, Xiao L, Lang W, Gao F, Ruvolo P, May WS Jr. Novel role for JNK as a stress-activated Bcl2 kinase. *J Biochem*. 2001;276(26):23681–8.
 61. Nowak G, Clifton GL, Godwin ML, Bakajsova D. Activation of ERK1/2 pathway mediates oxidant-induced decreases in mitochondrial function in renal cells. *Am J Physiol Renal Physiol*. 2006;291(4):F840–55.
 62. Monick MM, Powers LS, Barrett CW, Hinde S, Ashare A, Groskreutz DJ, Nyunoya T, Coleman M, Spitz DR, Hunninghake GW. Constitutive ERK MAPK activity regulates macrophage ATP production and mitochondrial integrity. *J Immunol*. 2008;180(11):7485–96.
 63. Dagda RK, Zhu J, Kulich SM, Chu CT. Mitochondrially localized ERK2 regulates mitophagy and autophagic cell stress: implications for Parkinson's disease. *Autophagy*. 2008;4(6):770–82.
 64. Zhuang S, Kinsey GR, Yan Y, Han J, Schnellmann RG. Extracellular signal-regulated kinase activation mediates mitochondrial dysfunction and necrosis induced by hydrogen peroxide in renal proximal tubular cells. *J Pharmacol Exp Ther*. 2008;325(3):732–40.
 65. Hernandez-Resendiz S, Zazueta C. PHO-ERK1/2 interaction with mitochondria regulates the permeability transition pore in cardioprotective signaling. *Life Sci*. 2014;108(1):13–21.
 66. Duarte A, Castillo AF, Podesta EJ, Poderoso C. Mitochondrial fusion and ERK activity regulate steroidogenic acute regulatory protein localization in mitochondria. *PLoS One*. 2014;9(6):e100387.
 67. Ballard-Croft C, Kristo G, Yoshimura Y, Reid E, Keith BJ, Mentzer RM Jr, Lasley RD. Acute adenosine preconditioning is mediated by p38 MAPK activation in discrete subcellular compartments. *Am J Physiol Heart Circ Physiol*. 2005;288(3):H1359–66.
 68. Galli S, Antico Arciuch VG, Poderoso C, Converso DP, Zhou Q, Bal de Kier Joffe E, Cadenas E, Boczkowski J, Carreras MC, Poderoso JJ. Tumor cell phenotype is sustained by selective MAPK oxidation in mitochondria. *PLoS One*. 2008;3(6):2379.
 69. Hoshi M, Sato M, Kondo S, Takashima A, Noguchi K, Takahashi M, Ishiguro K, Imahori K. Different localization of tau protein kinase I/glycogen synthase kinase-3 beta from glycogen synthase kinase-3 alpha in cerebellum mitochondria. *J Biochem*. 1995;118(4):683–5.
 70. Bijur GN, Jope RS. Glycogen synthase kinase-3 beta is highly activated in nuclei and mitochondria. *Neuro Rep*. 2003;14(18):2415–9.
 71. Nishihara M, Miura T, Miki T, Tanno M, Yano T, Naitoh K, Ohori K, Hotta H, Terashima Y, Shimamoto K. Modulation of the mitochondrial permeability transition pore complex in GSK-3beta-mediated myocardial protection. *J Mol Cell Cardiol*. 2007;43(5):564–70.
 72. Xi J, Wang H, Mueller RA, Norfleet EA, Xu Z. Mechanism for resveratrol-induced cardioprotection against reperfusion injury involves glycogen synthase kinase 3beta and mitochondrial permeability transition pore. *Eur J Pharmacol*. 2009;604(1–3):111–6.
 73. Rodriguez-Sinovas A, Boengler K, Cabestrero A, Gres P, Morente M, Ruiz-Meana M, Konietzka I, Miro E, Totzeck A, Heusch G, et al. Translocation of connexin 43 to the inner mitochondrial membrane of cardiomyocytes through the heat shock protein 90-dependent TOM pathway and its importance for cardioprotection. *Circ Res*. 2006;99(1):93–101.
 74. Hoshi M, Takashima A, Noguchi K, Murayama M, Sato M, Kondo S, Saitoh Y, Ishiguro K, Hoshino T, Imahori K. Regulation of mitochondrial pyruvate dehydrogenase activity by tau protein kinase I/glycogen synthase kinase 3beta in brain. *Proc Natl Acad Sci USA*. 1996;93(7):2719–23.
 75. Juhaszova M, Zorov DB, Kim SH, Pepe S, Fu Q, Fishbein KW, Ziman BD, Wang S, Ytrehus K, Antos CL, et al. Glycogen synthase kinase-3beta mediates convergence of protection signaling to inhibit the mitochondrial permeability transition pore. *J Clin Invest*. 2004;113(11):1535–49.
 76. Hanger DP, Hughes K, Woodgett JR, Brion JP, Anderton BH. Glycogen synthase kinase-3 induces Alzheimer's disease-like phosphorylation

- of tau: generation of paired helical filament epitopes and neuronal localisation of the kinase. *Neurosci Lett*. 1992;147(1):58–62.
77. Mulot SF, Hughes K, Woodgett JR, Anderton BH, Hanger DP. PHF-tau from Alzheimer's brain comprises four species on SDS-PAGE which can be mimicked by in vitro phosphorylation of human brain tau by glycogen synthase kinase-3 beta. *FEBS Lett*. 1994;349(3):359–64.
 78. Oliveira JM, Henriques AG, Martins F, Rebelo S, da Cruz E Silva OA. Amyloid-beta modulates both A β PP and tau phosphorylation. *J Alzheimer Dis*. 2015;45(2):495–507.
 79. Kwok JB, Hallupp M, Loy CT, Chan DK, Woo J, Mellick GD, Buchanan DD, Silburn PA, Halliday GM, Schofield PR. GSK3B polymorphisms alter transcription and splicing in Parkinson's disease. *Ann Neurol*. 2005;58(6):829–39.
 80. Ma C, Wang J, Gao Y, Gao TW, Chen G, Bower KA, Odetallah M, Ding M, Ke Z, Luo J. The role of glycogen synthase kinase 3beta in the transformation of epidermal cells. *Cancer Res*. 2007;67(16):7756–64.
 81. Farago M, Dominguez I, Landesman-Bollag E, Xu X, Rosner A, Cardiff RD, Seldin DC. Kinase-inactive glycogen synthase kinase 3beta promotes Wnt signaling and mammary tumorigenesis. *Cancer Res*. 2005;65(13):5792–801.
 82. Vincent T, Kukalev A, Andang M, Pettersson R, Percipalle P. The glycogen synthase kinase (GSK) 3beta represses RNA polymerase I transcription. *Oncogene*. 2008;27(39):5254–9.
 83. Shakoori A, Mai W, Miyashita K, Yasumoto K, Takahashi Y, Ooi A, Kawakami K, Minamoto T. Inhibition of GSK-3 beta activity attenuates proliferation of human colon cancer cells in rodents. *Cancer Sci*. 2007;98(9):1388–93.
 84. Ougolkov AV, Fernandez-Zapico ME, Savoy DN, Urrutia RA, Billadeau DD. Glycogen synthase kinase-3beta participates in nuclear factor kappaB-mediated gene transcription and cell survival in pancreatic cancer cells. *Cancer Res*. 2005;65(6):2076–81.
 85. Papa S, Sardanelli AM, Scacco S, Technikova-Dobrova Z. cAMP-dependent protein kinase and phosphoproteins in mammalian mitochondria. An extension of the cAMP-mediated intracellular signal transduction. *FEBS Lett*. 1999;444(2):245–9.
 86. Kleitke B, Sydow H, Wollenberger A. Evidence for cyclic AMP-dependent protein kinase activity in isolated guinea pig and rat liver mitochondria. *Acta Biologica et medica Germanica*. 1976;35(3–4):K9–17.
 87. Schwoch G, Trinczek B, Bode C. Localization of catalytic and regulatory subunits of cyclic AMP-dependent protein kinases in mitochondria from various rat tissues. *Biochem J*. 1990;270(1):181–8.
 88. Verdanis A. Protein kinase activity at the inner membrane of mammalian mitochondria. *J Biochem*. 1977;252(3):807–13.
 89. Ma MP, Thomson M. Protein Kinase A Subunit alpha Catalytic and A Kinase Anchoring Protein 79 in Human Placental Mitochondria. *Open Biochem J*. 2012;6:23–30.
 90. Sardanelli AM, Technikova-Dobrova Z, Speranza F, Mazzocca A, Scacco S, Papa S. Topology of the mitochondrial cAMP-dependent protein kinase and its substrates. *FEBS Lett*. 1996;396(2–3):276–8.
 91. Technikova-Dobrova Z, Sardanelli AM, Papa S. Phosphorylation of mitochondrial proteins in bovine heart. Characterization of kinases and substrates. *FEBS Lett*. 1993;322(1):51–5.
 92. Technikova-Dobrova Z, Sardanelli AM, Stanca MR, Papa S. cAMP-dependent protein phosphorylation in mitochondria of bovine heart. *FEBS Lett*. 1994;350(2–3):187–91.
 93. Gomez-Chang E, Espinosa-Garcia MT, Olvera-Sanchez S, Flores-Herrera O, Martinez F. PKA tightly bound to human placental mitochondria participates in steroidogenesis and is not modified by cAMP. *Placenta*. 2014;35(9):748–62.
 94. Feliciello A, Gottesman ME, Avvedimento EV. The biological functions of A-kinase anchor proteins. *J Mol Biol*. 2001;308(2):99–114.
 95. Tasken K, Aandahl EM. Localized effects of cAMP mediated by distinct routes of protein kinase A. *Physiol Rev*. 2004;84(1):137–67.
 96. Harada H, Becknell B, Wilm M, Mann M, Huang LJ, Taylor SS, Scott JD, Kormeyer SJ. Phosphorylation and inactivation of BAD by mitochondria-anchored protein kinase A. *Mol Cell*. 1999;3(4):413–22.
 97. Affaitati A, Cardone L, de Cristofaro T, Carlucci A, Ginsberg MD, Varone S, Gottesman ME, Avvedimento EV, Feliciello A. Essential role of A-kinase anchor protein 121 for cAMP signaling to mitochondria. *J Biochem*. 2003;278(6):4286–94.
 98. Merrill RA, Strack S. Mitochondria: a kinase anchoring protein 1, a signaling platform for mitochondrial form and function. *Int J Biochem Cell Biol*. 2014;48:92–6.
 99. Cribbs JT, Strack S. Reversible phosphorylation of Drp1 by cyclic AMP-dependent protein kinase and calcineurin regulates mitochondrial fission and cell death. *EMBO Rep*. 2007;8(10):939–44.
 100. Dickey AS, Strack S. PKA/AKAP1 and PP2A/Bbeta2 regulate neuronal morphogenesis via Drp1 phosphorylation and mitochondrial biogenesis. *J Neurosci*. 2011;31(44):15716–26.
 101. Lee J, Kim CH, Simon DK, Aminova LR, Andreyev AY, Kushnareva YE, Murphy AN, Lonze BE, Kim KS, Ginty DD, et al. Mitochondrial cyclic AMP response element-binding protein (CREB) mediates mitochondrial gene expression and neuronal survival. *J Biochem*. 2005;280(49):40398–401.
 102. Ryu H, Lee J, Impey S, Ratan RR, Ferrante RJ. Antioxidants modulate mitochondrial PKA and increase CREB binding to D-loop DNA of the mitochondrial genome in neurons. *Proc Nat Acad Sci USA*. 2005;102(39):13915–20.
 103. Beall A, Bagwell D, Woodrum D, Stoming TA, Kato K, Suzuki A, Rasmussen H, Brophy CM. The small heat shock-related protein, HSP20, is phosphorylated on serine 16 during cyclic nucleotide-dependent relaxation. *J Biochem*. 1999;274(16):11344–51.
 104. Edwards HV, Scott JD, Baillie GS. PKA phosphorylation of the small heat-shock protein Hsp20 enhances its cardioprotective effects. *Biochem Soc Trans*. 2012;40(1):210–4.
 105. Ohnuma Y, Miura T, Miki T, Tanno M, Kuno A, Tsuchida A, Shimamoto K. Opening of mitochondrial K(ATP) channel occurs downstream of PKC-epsilon activation in the mechanism of preconditioning. *Am J Physiol Heart Circ Physiol*. 2002;283(1):H440–7.
 106. Ogbi M, Johnson JA. Protein kinase Cepsilon interacts with cytochrome c oxidase subunit IV and enhances cytochrome c oxidase activity in neonatal cardiac myocyte preconditioning. *Biochem J*. 2006;393(Pt 1):191–9.
 107. Ping P, Song C, Zhang J, Guo Y, Cao X, Li RC, Wu W, Vondriska TM, Pass JM, Tang XL, et al. Formation of protein kinase C(epsilon)-Lck signaling modules confers cardioprotection. *J Clin Invest*. 2002;109(4):499–507.
 108. Jaburek M, Costa AD, Burton JR, Costa CL, Garlid KD. Mitochondrial PKC epsilon and mitochondrial ATP-sensitive K+ channel copurify and coreconstitute to form a functioning signaling module in proteoliposomes. *Circ Res*. 2006;99(8):878–83.
 109. Waza AA, Andrabi K, Hussain MU. Protein kinase C (PKC) mediated interaction between connexin43 (Cx43) and K(+)-ATP channel subunit (Kir6.1) in cardiomyocyte mitochondria: implications in cytoprotection against hypoxia induced cell apoptosis. *Cell Signal*. 2014;26(9):1909–17.
 110. Hamasaki Y, Shinohara O, Ishida H, Hayashi Y, Nakazawa H. Decreased protein kinase C-epsilon expression in hypertrophied cardiac ventricles induced by triiodothyronine treatment in the rat. *Life Sci*. 2000;67(15):1859–68.
 111. Inagaki K, Iwanaga Y, Sarai N, Onozawa Y, Takenaka H, Mochly-Rosen D, Kihara Y. Tissue angiotensin II during progression or ventricular hypertrophy to heart failure in hypertensive rats; differential effects on PKC epsilon and PKC beta. *J Mol Cell Cardiol*. 2002;34(10):1377–85.
 112. Chen L, Hahn H, Wu G, Chen X, Liron T, Schechtman D, Cavallaro G, Banci L, Guo Y, Bolli R, et al. Opposing cardioprotective actions and parallel hypertrophic effects of delta PKC and epsilon PKC. *Proc Nat Acad Sci USA*. 2001;98(20):11114–9.
 113. Choi DS, Wang D, Yu GQ, Zhu G, Kharaznia VN, Paredes JP, Chang WS, Deitchman JK, Mucke L, Messing RO. PKCepsilon increases endothelin converting enzyme activity and reduces amyloid plaque pathology in transgenic mice. *Proc Nat Acad Sci USA*. 2006;103(21):8215–20.
 114. Majumder PK, Pandey P, Sun X, Cheng K, Datta R, Saxena S, Kharbanda S, Kufe D. Mitochondrial translocation of protein kinase C delta in phorbol ester-induced cytochrome c release and apoptosis. *J Biochem*. 2000;275(29):21793–6.
 115. Majumder PK, Mishra NC, Sun X, Bharti A, Kharbanda S, Saxena S, Kufe D. Targeting of protein kinase C delta to mitochondria in the oxidative stress response. *Cell Grow Differ*. 2001;12(9):465–70.
 116. Sumitomo M, Ohba M, Asakuma J, Asano T, Kuroki T, Hayakawa M. Protein kinase Cdelta amplifies ceramide formation via mitochondrial signaling in prostate cancer cells. *J Clin Invest*. 2002;109(6):827–36.

117. Li L, Lorenzo PS, Bogi K, Blumberg PM, Yuspa SH. Protein kinase Cdelta targets mitochondria, alters mitochondrial membrane potential, and induces apoptosis in normal and neoplastic keratinocytes when over-expressed by an adenoviral vector. *Mol Cell Biol*. 1999;19(12):8547–58.
118. Valente EM, Abou-Sleiman PM, Caputo V, Muqit MM, Harvey K, Gispert S, Ali Z, Del Turco D, Bentivoglio AR, Healy DG, et al. Hereditary early-onset Parkinson's disease caused by mutations in PINK1. *Science*. 2004;304(5674):1158–60.
119. Weihofen A, Thomas KJ, Ostaszewski BL, Cookson MR, Selkoe DJ. Pink1 forms a multiprotein complex with Miro and Milton, linking Pink1 function to mitochondrial trafficking. *Biochemistry*. 2009;48(9):2045–52.
120. Jin SM, Lazarou M, Wang C, Kane LA, Narendra DP, Youle RJ. Mitochondrial membrane potential regulates PINK1 import and proteolytic destabilization by PARL. *J Cell Biol*. 2010;191(5):933–42.
121. Gasser T. Genetics of Parkinson's disease. *Dial Clin Neurosci*. 2004;6(3):295–301.
122. Chen Y, Sparks M, Bhandari P, Matkovich SJ, Dorn GW. Mitochondrial genome linearization is a causative factor for cardiomyopathy in mice and drosophila. *Antioxid Redox Signal*. 2013;21(4):1949–59.
123. Wang X, Winter D, Ashrafi G, Schlehe J, Wong YL, Selkoe D, Rice S, Steen J, LaVoie MJ, Schwarz TL. PINK1 and Parkin target Miro for phosphorylation and degradation to arrest mitochondrial motility. *Cell*. 2011;147(4):893–906.
124. Dagda RK, Cherra SJ 3rd, Kulich SM, Tandon A, Park D, Chu CT. Loss of PINK1 function promotes mitophagy through effects on oxidative stress and mitochondrial fission. *J Biochem*. 2009;284(20):13843–55.
125. Lutz AK, Exner N, Fett ME, Schlehe JS, Kloos K, Lammermann K, Brunner B, Kurz-Drexler A, Vogel F, Reichert AS, et al. Loss of parkin or PINK1 function increases Drp1-dependent mitochondrial fragmentation. *J Biochem*. 2009;284(34):22938–51.
126. Liu S, Sawada T, Lee S, Yu W, Silverio G, Alapatt P, Millan I, Shen A, Saxton W, Kanao T, et al. Parkinson's disease-associated kinase PINK1 regulates Miro protein level and axonal transport of mitochondria. *PLoS Genet*. 2012;8(3):e1002537.
127. Gautier CA, Kitada T, Shen J. Loss of PINK1 causes mitochondrial functional defects and increased sensitivity to oxidative stress. *Proc Nat Acad Sci USA*. 2008;105(32):11364–9.
128. Morais VA, Verstreken P, Roethig A, Smet J, Snellinx A, Vanbrabant M, Haddad D, Frezza C, Mandemakers W, Vogt-Weisenhorn D, et al. Parkinson's disease mutations in PINK1 result in decreased Complex I activity and deficient synaptic function. *EMBO Mol Med*. 2009;1(2):99–111.
129. Vos M, Esposito G, Edirisinghe JN, Vilain S, Haddad DM, Slabbaert JR, Van Meensel S, Schaap O, De Strooper B, Meganathan R, et al. Vitamin K2 is a mitochondrial electron carrier that rescues pink1 deficiency. *Science*. 2012;336(6086):1306–10.
130. Vincow ES, Merrihew G, Thomas RE, Shulman NJ, Beyer RP, MacCoss MJ, Pallanck LJ. The PINK1-Parkin pathway promotes both mitophagy and selective respiratory chain turnover in vivo. *Proc Nat Acad Sci USA*. 2013;110(16):6400–5.
131. Gehrke S, Wu Z, Klinkenberg M, Sun Y, Auburger G, Guo S, Lu B. PINK1 and Parkin control localized translation of respiratory chain component mRNAs on mitochondria outer membrane. *Cell Metab*. 2015;21(1):95–108.
132. Sekine S, Kanamaru Y, Koike M, Nishihara A, Okada M, Kinoshita H, Kamiyama M, Maruyama J, Uchiyama Y, Ishihara N, et al. Rhomboid protease PARL mediates the mitochondrial membrane potential loss-induced cleavage of PGAM5. *J Biochem*. 2012;287(41):34635–45.
133. Moiso N, Fedele V, Edwards J, Martins LM. Loss of PINK1 enhances neurodegeneration in a mouse model of Parkinson's disease triggered by mitochondrial stress. *Neuropharmacology*. 2014;77:350–7.
134. Tan EK, Refai FS, Siddique M, Yap K, Ho P, Fook-Chong S, Zhao Y. Clinically reported heterozygous mutations in the PINK1 kinase domain exert a gene dosage effect. *Hum Mutat*. 2009;30(11):1551–7.
135. Sim CH, Lio DS, Mok SS, Masters CL, Hill AF, Culvenor JG, Cheng HC. C-terminal truncation and Parkinson's disease-associated mutations down-regulate the protein serine/threonine kinase activity of PTEN-induced kinase-1. *Hum Mol Genet*. 2006;15(21):3251–62.
136. Inzelberg R, Jankovic J. Are Parkinson disease patients protected from some but not all cancers? *Neurology*. 2007;69(15):1542–50.
137. Siddall HK, Warrell CE, Davidson SM, Mocanu MM, Yellon DM. Mitochondrial PINK1—a novel cardioprotective kinase? *Cardiovasc Drug*. 2008;22(6):507–8.
138. Scheele C, Nielsen AR, Walden TB, Sewell DA, Fischer CP, Brogan RJ, Petrovic N, Larsson O, Tesch PA, Wennmalm K, et al. Altered regulation of the PINK1 locus: a link between type 2 diabetes and neurodegeneration? *FASEB J*. 2007;21(13):3653–65.
139. Gandhi S, Wood-Kaczmar A, Yao Z, Plun-Favreau H, Deas E, Klupsch K, Downward J, Latchman DS, Tabrizi SJ, Wood NW, et al. PINK1-associated Parkinson's disease is caused by neuronal vulnerability to calcium-induced cell death. *Mol Cell*. 2009;33(5):627–38.
140. Deas E, Piipari K, Machhada A, Li A, Gutierrez-del-Arroyo A, Withers DJ, Wood NW, Abramov AY. PINK1 deficiency in beta-cells increases basal insulin secretion and improves glucose tolerance in mice. *Open Biol*. 2014;4:140051.
141. Wu JJ, Zhang L, Bennett AM. The noncatalytic amino terminus of mitogen-activated protein kinase phosphatase 1 directs nuclear targeting and serum response element transcriptional regulation. *Mol Cell Biol*. 2005;25(11):4792–803.
142. Rosini P, De Chiara G, Bonini P, Lucibello M, Marcocci ME, Garaci E, Cozzolino F, Torcia M. Nerve growth factor-dependent survival of CESS B cell line is mediated by increased expression and decreased degradation of MAPK phosphatase 1. *J Biochem*. 2004;279(14):14016–23.
143. Wang HY, Cheng Z, Malbon CC. Overexpression of mitogen-activated protein kinase phosphatases MKP1, MKP2 in human breast cancer. *Cancer Lett*. 2003;191(2):229–37.
144. Loda M, Capodiceci P, Mishra R, Yao H, Corless C, Grigioni W, Wang Y, Magi-Galluzzi C, Stork PJ. Expression of mitogen-activated protein kinase phosphatase-1 in the early phases of human epithelial carcinogenesis. *Am J Pathol*. 1996;149(5):1553–64.
145. Denkert C, Schmitt WD, Berger S, Reles A, Pest S, Siegert A, Lichtenegger W, Dietel M, Hauptmann S. Expression of mitogen-activated protein kinase phosphatase-1 (MKP-1) in primary human ovarian carcinoma. *Int J Cancer*. 2002;102(5):507–13.
146. Chattopadhyay S, Machado-Pinilla R, Manguan-Garcia C, Belda-Iniesta C, Moratilla C, Cejas P, Fresno-Vara JA, de Castro-Carpeno J, Casado E, Nistal M, et al. MKP1/CL100 controls tumor growth and sensitivity to cisplatin in non-small-cell lung cancer. *Oncogene*. 2006;25(23):3335–45.
147. Moncho-Amor V, de Caceres II, Bandres E, Martinez-Poveda B, Orgaz JL, Sanchez-Perez I, Zazo S, Rovira A, Albanell J, Jimenez B, et al. DUSP1/MKP1 promotes angiogenesis, invasion and metastasis in non-small-cell lung cancer. *Oncogene*. 2011;30(6):668–78.
148. Salojin KV, Owusu IB, Millerchip KA, Potter M, Platt KA, Oravec T. Essential role of MAPK phosphatase-1 in the negative control of innate immune responses. *J Immunol*. 2006;176(3):1899–907.
149. Chi H, Barry SP, Roth RJ, Wu JJ, Jones EA, Bennett AM, Flavell RA. Dynamic regulation of pro- and anti-inflammatory cytokines by MAPK phosphatase 1 (MKP-1) in innate immune responses. *Proc Nat Acad Sci USA*. 2006;103(7):2274–9.
150. Zhao Q, Shepherd EG, Manson ME, Nelin LD, Sorokin A, Liu Y. The role of mitogen-activated protein kinase phosphatase-1 in the response of alveolar macrophages to lipopolysaccharide: attenuation of proinflammatory cytokine biosynthesis via feedback control of p38. *J Biochem*. 2005;280(9):8101–8.
151. Salvi M, Stringaro A, Brunati AM, Agostinelli E, Arancia G, Clari G, Toninello A. Tyrosine phosphatase activity in mitochondria: presence of Shp-2 phosphatase in mitochondria. *Cell Mol Life Sci*. 2004;61(18):2393–404.
152. Zang QS, Martinez B, Yao X, Maass DL, Ma L, Wolf SE, Minei JP. Sepsis-induced cardiac mitochondrial dysfunction involves altered mitochondrial-localization of tyrosine kinase Src and tyrosine phosphatase SHP2. *PLoS One*. 2012;7(8):e43424.
153. Tartaglia M, Mehler EL, Goldberg R, Zampino G, Brunner HG, Kremer H, van der Burg I, Crosby AH, Ion A, Jeffery S, et al. Mutations in PTPN11, encoding the protein tyrosine phosphatase SHP-2, cause Noonan syndrome. *Nature Genet*. 2001;29(4):465–8.
154. Legius E, Schrandt-Stumpel C, Schollen E, Pulles-Heintzberger C, Gewillig M, Fryns JP. PTPN11 mutations in LEOPARD syndrome. *J Med Genet*. 2002;39(8):571–4.

155. Tartaglia M, Gelb BD. Noonan syndrome and related disorders: genetics and pathogenesis. *Ann Rev Genomics Hum Genet.* 2005;6:45–68.
156. Kontaridis MI, Swanson KD, David FS, Barford D, Neel BG. PTPN11 (Shp2) mutations in LEOPARD syndrome have dominant negative, not activating, effects. *J Biochem.* 2006;281(10):6785–92.
157. Loh ML, Vattikuti S, Schubbert S, Reynolds MG, Carlson E, Lieu KH, Cheng JW, Lee CM, Stokoe D, Bonifas JM, et al. Mutations in PTPN11 implicate the SHP-2 phosphatase in leukemogenesis. *Blood.* 2004;103(6):2325–31.
158. Emanuel PD, Shannon KM, Castleberry RP. Juvenile myelomonocytic leukemia: molecular understanding and prospects for therapy. *Mol Med Today.* 1996;2(11):468–75.
159. Merks JH, Caron HN, Hennekam RC. High incidence of malformation syndromes in a series of 1073 children with cancer. *Am J Med Genet Part A.* 2005;134A(2):132–43.
160. Ucar C, Calyskan U, Martini S, Heinritz W. Acute myelomonocytic leukemia in a boy with LEOPARD syndrome (PTPN11 gene mutation positive). *J Pediatr Hematol Oncol.* 2006;28(3):123–5.
161. Chatelain EH, Dupuy JW, Letellier T, Dachary-Prigent J. Functional impact of PTP1B-mediated Src regulation on oxidative phosphorylation in rat brain mitochondria. *Cell Mol Life Sci.* 2011;68(15):2603–13.
162. Pagliarini DJ, Wiley SE, Kimple ME, Dixon JR, Kelly P, Worby CA, Casey PJ, Dixon JE. Involvement of a mitochondrial phosphatase in the regulation of ATP production and insulin secretion in pancreatic beta cells. *Mol Cell.* 2005;19(2):197–207.
163. Vazquez F, Matsuoka S, Sellers WR, Yanagida T, Ueda M, Devreotes PN. Tumor suppressor PTEN acts through dynamic interaction with the plasma membrane. *Proc Natl Acad Sci USA.* 2006;103(10):3633–8.
164. Zhu Y, Hoell P, Ahlemeyer B, Kriegelstein J. PTEN: a crucial mediator of mitochondria-dependent apoptosis. *Apoptosis.* 2006;11(2):197–207.
165. Zu L, Zheng X, Wang B, Parajuli N, Steenbergen C, Becker LC, Cai ZP. Ischemic preconditioning attenuates mitochondrial localization of PTEN induced by ischemia-reperfusion. *Am J Physiol Heart Circ Physiol.* 2011;300(6):H2177–86.
166. Liang H, He S, Yang J, Jia X, Wang P, Chen X, Zhang Z, Zou X, McNutt MA, Shen WH, et al. PTENalpha, a PTEN isoform translated through alternative initiation, regulates mitochondrial function and energy metabolism. *Cell Metab.* 2014;19(5):836–48.
167. Endo T, Kohda D. Functions of outer membrane receptors in mitochondrial protein import. *Biochim Biophys Acta.* 2002;1592(1):3–14.
168. Meisinger C, Ryan MT, Hill K, Model K, Lim JH, Sickmann A, Muller H, Meyer HE, Wagner R, Pfanner N. Protein import channel of the outer mitochondrial membrane: a highly stable Tom40-Tom22 core structure differentially interacts with preproteins, small tom proteins, and import receptors. *Mol Cell Biol.* 2001;21(7):2337–48.
169. Wiedemann N, Kozjak V, Chacinska A, Schonfisch B, Rospert S, Ryan MT, Pfanner N, Meisinger C. Machinery for protein sorting and assembly in the mitochondrial outer membrane. *Nature.* 2003;424(6948):565–71.
170. Rehling P, Model K, Brandner K, Kovermann P, Sickmann A, Meyer HE, Kuhlbrandt W, Wagner R, Truscott KN, Pfanner N. Protein insertion into the mitochondrial inner membrane by a twin-pore translocase. *Science.* 2003;299(5613):1747–51.
171. Mesecke N, Terziyska N, Kozany C, Baumann F, Neupert W, Hell K, Herrmann JM. A disulfide relay system in the intermembrane space of mitochondria that mediates protein import. *Cell.* 2005;121(7):1059–69.
172. Covian R, Balaban RS. Cardiac mitochondrial matrix and respiratory complex protein phosphorylation. *Am J Physiol Heart Circ Physiol.* 2012;303(8):H940–66.
173. Feng J, Zhu M, Schaub MC, Gehrig P, Roschitzki B, Lucchinetti E, Zaugg M. Phosphoproteome analysis of isoflurane-protected heart mitochondria: phosphorylation of adenine nucleotide translocator-1 on Tyr194 regulates mitochondrial function. *Cardiovas Res.* 2008;80(1):20–9.
174. Grimsrud PA, Carson JJ, Hebert AS, Hubler SL, Niemi NM, Bailey DJ, Jochem A, Stapleton DS, Keller MP, Westphall MS, et al. A quantitative map of the liver mitochondrial phosphoproteome reveals posttranslational control of ketogenesis. *Cell Metab.* 2012;16(5):672–83.
175. Pagliarini DJ, Dixon JE. Mitochondrial modulation: reversible phosphorylation takes center stage? *Trend Biochem Sci.* 2006;31(1):26–34.
176. Zhao X, Leon IR, Bak S, Mogensen M, Wrzesinski K, Hojlund K, Jensen ON. Phosphoproteome analysis of functional mitochondria isolated from resting human muscle reveals extensive phosphorylation of inner membrane protein complexes and enzymes. *Mol Cell Proteom.* 2011;10(1):110000299.

Submit your next manuscript to BioMed Central
and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

